
What is a set?

Leeds Logic Seminar

Leeds, November 24, 2010

.

Peter Aczel

`petera@cs.man.ac.uk`

Manchester University,

Plan of the talk

- Introduction
- The classical conception of set
- The type theoretic conception of set
- A quick look at the type theory
- Some references

Some set theories

$$\begin{array}{ccccc} CZF & \Rightarrow & IZF & \Rightarrow & ZF \\ \uparrow & & \uparrow & & \Downarrow \\ CZF_{R,E} & \Rightarrow & IZF_R & & (CZF_{R,E} + EM) \end{array}$$

$$ZF \leq_{\neg\neg} IZF$$

$$CZF \sim ID_1 \sim KP \ll IZF \sim ZF$$

The type-theoretic interpretations

$$CZF \leq_{tt} (ML + \mathbb{U} + \mathbb{V})$$

$$ZF \leq_{tt} (ML + \mathbb{U} + \mathbb{V} + EM)$$

$$ZF \leq_{\neg\neg} IZF \leq_{tt} (ML + \mathbb{Q} + \mathbb{U} + \mathbb{V})$$

The cumulative hierarchy in ZF

- $V = \bigcup_{\alpha \in On} V_\alpha$,
where $V_\alpha = \bigcup_{\beta < \alpha} Pow(V_\beta)$ for $\alpha \in On$.
- $V_0 \subseteq V_1 \subseteq \dots V_\alpha \subseteq V_{\alpha+1} \subseteq \dots$
- $V_0 = \emptyset, V_1 = \{\emptyset\}, V_{\alpha+1} = Pow(V_\alpha)$
- $V_\alpha =$ the **set** of sets formed by stage α
- $\{x \in V_\beta \mid \dots x \dots\} \in V_\alpha$ if $\beta < \alpha$

The classical iterative conception

Zermelo, Scott, Schoenfield, Boolos, Parsons,...

- Sets are extensional.
- Sets are formed in stages, α , out of elements formed at earlier stages.
- A set is formed by collecting together (into a whole) its elements.
- There are lots of stages:
 1. There is a stage.
 2. For each stage there is a later stage.
 3. There is a stage, ω , reflecting 1,2.
 4. If $\{\alpha_i\}_{i \in I}$ is a family of stages indexed by a set I then there is a stage later than all the α_i .

The formation of powersets

Suppose X is an infinite set formed at some stage α .

- Then each element of X will have been formed at some stage before α .
- So each subset of X will have been formed at or before stage α . But can the elements of each subset of X , however the subset is formed, really be collected into a whole at or before stage α ?
- So the powerset, $Pow(X)$, will be formed at any stage after α . But can the subsets of X be collected into a whole so easily?

Types, sets and classes

- Any discussion concerning the concept of set must distinguish between the three distinct notions of **type**, **set** and **class**.
- For example the intended universe of ZF set theory is a **type**, the objects in that universe are **sets** and $\{x \mid x \notin x\}$ is a **class**.
- The notion of **type** is perhaps best taken as a pre-mathematical philosophical notion.
- The notion of **set** of ZF is an iterative combinatorial notion.
- The notion of **class** is a logical notion - the extension of a predicate.

Types and Classes

- A mathematical **object** is always given as an object of some **type**.
- We write $a : A$ for the **judgement** that a is an object of type A .
- A **class** on a type is the extension of a **propositional function** on the type.
- If B is a propositional function on the type A then its **extension** is the class $C = \{x : A \mid B(x)\}$.
- For $a : A$ the **proposition** that a is in the class C is $B(a)$.
- If also $C' = \{x : A \mid B'(x)\}$ then $(C = C')$ is the proposition

$$(\forall x : A)[B(x) \leftrightarrow B'(x)].$$

What is a **set of elements** of a type?

It is a **collection** into a whole of objects **chosen** from the type.

- e.g. given the type \mathbb{N} of natural numbers we have sets of natural numbers such as $\{0\}$, $\{0, 1\}$, $\{0, 3, 18\}$, $\{\}$
- and sets $\{0, 2, 4, 6, \dots, 92\}$, $\{2i \mid i < n\}$ for $n : \mathbb{N}$,
- and infinite sets such as $\{0, 2, 4, 6, \dots\} = \{2i \mid i : \mathbb{N}\}$.
- In general we can form sets of natural numbers $\{a_i \mid i \in I\}$ with $a_i : \mathbb{N}$ for $i : I$, where I is an **index-type**.

Index-types

- I use the word **index-type** for something like
 - Bishop's constructive notion of set,which I think is also something like
 - Martin-Löf's type-theoretic notion of set or data-typeand something like
 - the category theorists' notion of setwhen they talk about a category of sets.
- I need a distinct word in order to avoid confusion with the combinatorial notion of set, which is what axiomatic set theory is about.
- A **set** is formed out of its elements. But an **index-type** is an object that is conceptually prior to its elements.
- The index-types form a type \mathbb{U} .

Sets, of elements of a type, 1

- Given a type A , a **set of elements of A** is given by:
 1. an index-type I , the **index-type** of the set,
 2. a function $f : I \rightarrow A$, also thought of as a **family of elements of A** , $\{a_i\}_{i:I}$, where $a_i = f(i) : A$ for $i : I$.
- We may write the set as $(\text{set } i : I) f(i)$ or $[a_i \mid i : I]$.
- The **chosen elements** of the set are the a_i for $i : I$.
- The sets of elements of A form a type $Sub(A)$.
- A itself may be an index-type, so that we may form the set $[x \mid x : A] : Sub(A)$ of all the objects of A .
- But, in general, the type A need not be an index-type.

Sets, of elements of a type, 2

- An **equality relation**, $=_A$, on a type A is an assignment of a proposition $(b =_A c)$ to $b, c : A$ so that the laws for an equivalence relation hold; i.e.
 - $(\forall x : A)[x =_A x]$, $(\forall x, y : A)[x =_A y \rightarrow y =_A x]$,
 - $(\forall x, y, z : A)[x =_A y \rightarrow (y =_A z \rightarrow x =_A z)]$.
- Given an equality relation $=_A$ on a type A we may define the **membership relation** \in_A and **extensional equality relation** $=_{Sub(A)}$ as follows:
 - If $\alpha : Sub(A)$ is $[a_i \mid i : I]$ then, for $a : A$, $(a \in_A \alpha)$ is the proposition $(\exists i : I)[a =_A a_i]$.
 - If also $\beta : Sub(A)$ is $[b_j \mid j : J]$ then $(\alpha =_{Sub(A)} \beta)$ is the proposition
$$(\forall i : I)(\exists j : J)[a_i =_A b_j] \wedge (\forall j : J)(\exists i : I)[a_i =_A b_j].$$

The type of iterative sets

- The type \mathbb{V} of iterative sets is the inductive type obtained by iterating the **set-of** operation.
- The iterative sets are generated using the following rule.

Any set-of objects in \mathbb{V} is an object in \mathbb{V}

- In Constructive Type Theory \mathbb{V} is the inductive type having the introduction rule

$$\frac{I : \mathbb{U} \quad f : I \rightarrow \mathbb{V}}{(\text{set } i : I) f(i) : \mathbb{V}}$$

- So we have $Sub(\mathbb{V}) = \mathbb{V}$.

Equality and membership on \mathbb{V}

- We can recursively define $(\alpha =_{\mathbb{V}} \beta)$ for $\alpha, \beta : V$ using the rule

$$\frac{(\forall i : I)(\exists j : J)[a_i =_{\mathbb{V}} b_j] \wedge (\forall j : J)(\exists i : I)[a_i =_{\mathbb{V}} b_j]}{\alpha =_{\mathbb{V}} \beta}$$

where $\alpha = [a_i \mid i : I]$ and $\beta = [b_j \mid j : J]$.

- Also

$$\alpha \in_V \beta := (\exists j : J)(\alpha =_{\mathbb{V}} b_j).$$

ZF justified in type theory

- The logic of Type theory is obtained using the **Propositions-as-types** paradigm.
- The logic using this paradigm is intuitionistic logic.
- Also the type-theoretic axiom of choice holds:

$$\begin{aligned} & (\forall x : A)(\exists y : B(x))R(x, y) \\ & \quad \rightarrow (\exists f : (\prod x : A)B(x))(\forall x : A)R(x, f(x)) \end{aligned}$$

- The type \mathbb{V} of iterative sets provides an interpretation of *CZF* (Constructive *ZF*).
- **If the logic of type theory is made classical** then \mathbb{V} provides an interpretation of *ZF*.
- But this last step is not constructive.

The basic type theory $ML, 1$

The primitive forms of type

$\mathbb{O}, \mathbb{1}, \mathbb{B}, \mathbb{N}$ type

$$\frac{A \text{ type} \quad B(x) \text{ type } (x:A)}{(\Pi x:A)B(x), (\Sigma x:A)B(x) \text{ type}}$$

The introduction rules

$\star : \mathbb{1}$

$\text{t}, \text{f} : \mathbb{B}$

$$\frac{b(x):B(x) \quad (x:A)}{(\lambda x:A)b(x):(\Pi x:A)B(x)}$$

$0 : \mathbb{N}$

$$\frac{n:\mathbb{N}}{\text{succ}(n):\mathbb{N}}$$

$$\frac{a:A \quad b:B(a)}{\text{pair}(a,b):(\Sigma x:A)B(x)}$$

Defined types

$$(A_1 \rightarrow A_2) := (\Pi_ : A_1)A_2$$

$$(A_1 \times A_2) := (\Sigma_ : A_1)A_2$$

No type dependency so far!

The basic type theory $ML, 2$

There are standard elimination rules for each form of type
e.g. the standard elimination rule for \mathbb{B} is

$$\frac{A(x) \text{ type } (x:\mathbb{B}) \quad c:\mathbb{B} \quad a_1:A(\mathfrak{t}) \quad a_2:A(\mathfrak{f})}{\left\{ \begin{array}{l} \text{cases}(c, a_1, a_2) : A(c) \\ \text{cases}(\mathfrak{t}, a_1, a_2) = a_1 : A(\mathfrak{t}) \\ \text{cases}(\mathfrak{f}, a_1, a_2) = a_2 : A(\mathfrak{f}) \end{array} \right.}$$

In order to have dependent types we also use

$$\frac{c:\mathbb{B} \quad A_1, A_2 \text{ type}}{\left\{ \begin{array}{l} \text{Cases}(c, A_1, A_2) \text{ type} \\ \text{Cases}(\mathfrak{t}, A_1, A_2) = A_1 \\ \text{Cases}(\mathfrak{f}, A_1, A_2) = A_2 \end{array} \right.}$$

The basic type theory $ML, 3$

More defined types

$$\begin{aligned}(A_1 + A_2) &:= (\Sigma x : \mathbb{B}) \text{Cases}(x, A_1, A_2) && (A_1, A_2 \text{ type}) \\ \text{True}(c) &:= \text{Cases}(c, \mathbb{1}, \mathbb{0}) && (c : \mathbb{B})\end{aligned}$$

So we have the derived introduction rules

$$\frac{a:A_1}{\text{pair}(\mathfrak{t}, a):A_1+A_2} \qquad \frac{a:A_2}{\text{pair}(\mathfrak{f}, a):A_1+A_2}$$

and the equalities

$$\text{True}(\mathfrak{t}) = \mathbb{1} \qquad \text{True}(\mathfrak{f}) = \mathbb{0}$$

The basic type theory $ML, 4$

Propositions-as-Types (à la Curry-Howard)
Proposition = Type

<i>Prop</i>	\perp	\top	$A_1 \supset A_2$	$A_1 \wedge A_2$	$A_1 \vee A_2$
<i>Type</i>	$\mathbb{0}$	$\mathbb{1}$	$A_1 \rightarrow A_2$	$A_1 \times A_2$	$A_1 + A_2$

<i>Prop</i>	$(\forall x : A)B(x)$	$(\exists x : A)B(x)$
<i>Type</i>	$(\Pi x : A)B(x)$	$(\Sigma x : A)B(x)$

Adding Excluded Middle

$$\frac{A \text{ type}}{\sigma(A) : (A + (A \rightarrow \mathbb{O}))}$$

or equivalently

$$\frac{A \text{ type}}{\left\{ \begin{array}{l} \tau(A) : \mathbb{B} \\ t(A) : (\text{True}(\tau(A)) \leftrightarrow A) \end{array} \right.}$$

Note: $(B \leftrightarrow A) := (B \rightarrow A) \times (A \rightarrow B)$

Adding a type Ω

$$\Omega \text{ type} \quad \frac{a : \Omega}{\mathbb{T}(a) \text{ type}} \quad \frac{A \text{ type}}{\left\{ \begin{array}{l} \tau(A) : \Omega \\ t(A) : (\mathbb{T}(\tau(A)) \leftrightarrow A) \end{array} \right.}$$

- Think of Ω as a type of **predicative propositions**, or perhaps **truth values**. $\mathbb{T}(a)$ expresses that A is true. These rules are analogous to Bertrand Russell's Axiom of Reducibility and express that each proposition is logically equivalent to a predicative proposition.
- Excluded Middle is the extreme version of Ω where $\Omega = \mathbb{B}$ and $\mathbb{T}(a) = \text{True}(a)$.

Adding a type universe \mathbb{U}

- The rules of \mathbb{U} express that \mathbb{U} is a type of types that reflects the previous type forming rules.
- Reflecting the rules of ML we get

$$\mathbb{U} \text{ type} \quad \frac{A:\mathbb{U}}{A \text{ type}} \quad \mathbb{O}, \mathbb{1}, \mathbb{B}, \mathbb{N} : \mathbb{U} \quad \frac{A:\mathbb{U} \quad B(x):\mathbb{U} (x:A)}{(\prod x:A)B(x), (\sum x:A)B(x):\mathbb{U}}$$

- Reflecting the \mathbb{Q} rules we get

$$\mathbb{Q} : \mathbb{U} \quad \frac{a:\mathbb{Q}}{\mathbb{T}(a):\mathbb{U}}$$

The type-theoretic interpretations

$$CZF \leq_{tt} (ML + \mathbb{U} + \mathbb{V})$$

$$ZF \leq_{tt} (ML + \mathbb{U} + \mathbb{V} + EM)$$

$$ZF \leq_{\neg\neg} IZF \leq_{tt} (ML + \mathbb{Q} + \mathbb{U} + \mathbb{V})$$

Martin-Löf's Philosophy

- Notes on Constructive Mathematics, 1970.
- An intuitionistic theory of types: predicative part, in Logic Colloquium '73, published 1975.
- Intuitionistic Type Theory, 1980 lectures, notes by Giovanni Sambin, published as a Bibliopolis book in 1984.
- On the meanings of the logical constants and the justifications of the logical laws, 1983 lectures, published in Nordic Journal of Philosophical Logic in 1996

The type-theoretic interpretation of CST

- Aczel, 1978,
choice principles:1982,
inductive definitions:1986
- Aczel and Rathjen, [Notes on CST](#), Mittag-Leffler report,
2000/2001
- Gambino and Aczel, [The generalised type-theoretic interpretation of CST](#), JSL, vol 71 (2006), pp. 67-103.