

Bryophytes: Oback Burn ND451866. 8 March 2010.

John Crossley and Rosemary McCance

- Calypogeia fissa (Common Pouchwort, a Liverwort)
- Mnium hornum (Swan's-neck Thyme-moss)
- Kindbergia praelonga (Common feather moss)
- Pseudotaxiphyllum elegans (Elegant Silk-moss)
- Thuidium tamariscinum (Common Tamarisk-moss)
- Pseudoscleropodium purum (Neat Feather-moss)
- Hylocomium splendens (Glittering Wood-moss)
- Ulota bruchii (Bruch's Pincushion)
- Ulota phyllantha (Frizzled Pincushion)
- Plagiomnium undulatum (Hart's-tongue Thyme-moss)
- Hypnum jutlandicum (Heath Plait-moss)
- Preissia quadrata (Narrow Mushroom-headed Liverwort)
 - a liverwort with a mainly northerly and westerly
 - distribution in the UK, a lime-lover
- Orthotrichum pulchellum (Elegant Bristle-moss)
 - a moss of scattered distribution in the UK
- Pellia endivifolia (Endive Pellia, a Liverwort)
- Rhytidiadelphus squarrosus (Springy Turf-moss)
- Rhytidiadelphus triquetrus (Big Shaggy-moss)
- Brachythecium rutabulum (Rough-stalked Feather-moss)
- Calliergonella cuspidata (Pointed Spear-moss)